

Perspective on Clinically-Relevant Antimicrobial Resistant Enterobacterales in Food: Closing the Gaps using Genomics

Constanza P. Diaz¹, Francisca Alvarez¹, Jose M. Munita², Sandra Cortés³, Andrea I. Moreno Switt^{3*}

¹Andres Bello University, Chile, ²Clínica Alemana, Universidad del Desarrollo, Chile, ³Pontificia Universidad Católica de Chile, Chile

Submitted to Journal:
Frontiers in Sustainable Food Systems

Specialty Section:
Agro-Food Safety

Article type:
Perspective Article

Manuscript ID:
667504

Received on:
13 Feb 2021

Revised on:
02 Apr 2021

Journal website link:
www.frontiersin.org

Conflict of interest statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest

Author contribution statement

CDG, FPA, AIMS wrote the manuscript. JMM, SC critically reviewed the manuscript.

Keywords

antimicrobial resistance, Genomics, clinically relevant bacteria, E coli, Klebsiella pneumoniae, Food Safety Resistant Bacteria in Food

Abstract

Word count: 208

Antimicrobial resistance is one of the most important public health concerns—it causes 700,000 deaths annually according to the World Health Organization (WHO). Enterobacterales such as *E. coli* and *Klebsiella pneumoniae*, have become resistant to many relevant antimicrobials including carbapenems and extended spectrum cephalosporins. These clinically relevant resistant Enterobacterales (CRRE) members are now globally distributed in the environment including different food types (meats, produce, dairy). Unlike known foodborne pathogens, CRRE are not usually part of most food surveillance systems. However, numerous reports of CRRE highlight the importance of these bacteria in food and have been shown to contribute to the overall crisis of antimicrobial resistance. This is especially important in the context of carriage of these pathogens by immuno-compromised individuals. CRRE infections upon consumption of contaminated food could colonize the human gastrointestinal tract and eventually be a source of systemic infections such as urinary tract infections or septicemia. While different aspects need to be considered to elucidate this, whole genome sequencing along with metadata could be used to understand genomic relationships of CRRE obtained from foods and humans, including isolates from clinical infections. Once robust scientific data is available on the role of CRRE in food, countries could move forward to better survey and control CRRE in food.

Contribution to the field

Enterobacterales as *E. coli* and *Klebsiella pneumoniae* resistant to relevant antimicrobials as carbapenems and extended spectrum cephalosporins are considered clinically relevant resistant Enterobacterales (CRRE), which are also globally distributed in the environment including different food types (meats, produce, dairy). Unlike known foodborne pathogens, CRRE are not commonly part of most food surveillance systems. However, numerous reports of CRRE highlight the importance of the presence of these bacteria in food in the overall crisis of antimicrobial resistance. Though, the public health impact of the presence of CRRE in food remains unknown. A plausible outcome of CRRE upon consumption of contaminated food is to colonize the human gastrointestinal tract and eventually be involved in urinary tract infections or bloodstream infections. The article represents a perspective on the use of genomics to better understand the role of CRRE in food.

Funding statement

This work was funded by FONDECYT 1181167 and by the ANID Millennium Science Initiative/ Millennium Initiative for Collaborative Research on Bacterial Resistance, MICROB-R, NCN17_081.

Ethics statements

Studies involving animal subjects

Generated Statement: No animal studies are presented in this manuscript.

Studies involving human subjects

Generated Statement: No human studies are presented in this manuscript.

Inclusion of identifiable human data

Generated Statement: No potentially identifiable human images or data is presented in this study.

In review

Data availability statement

Generated Statement: The original contributions presented in the study are included in the article/supplementary material, further inquiries can be directed to the corresponding author/s.

In review

Perspective on Clinically-Relevant Antimicrobial Resistant Enterobacterales in Food: Closing the Gaps using Genomics

1

2 **Constanza Díaz-Gavidia^{1,2}, Francisca P. Álvarez^{1,2}, Jose M. Munita^{2,3}, Sandra Cortés^{4,5}, Andrea**
3 **I. Moreno-Switt^{2,6*}**

4 ¹ Facultad de Ciencias de la Vida, Universidad Andres Bello, Santiago, Chile

5 ² Millennium Initiative for Collaborative Research on Bacterial Resistance (MICROB-R), Santiago,
6 Chile

7 ³ Genomics and Resistant Microbes Group, Facultad de Medicina Clínica Alemana, Universidad del
8 Desarrollo, Santiago, Chile

9 ⁴ Escuela de Medicina, Advanced Center for Chronic Diseases (ACCDiS), Pontificia Universidad
10 Católica de Chile, Santiago, Chile

11 ⁵ Centro de Desarrollo Urbano Sustentable, Pontificia Universidad Católica de Chile.

12 ⁶ Escuela de Medicina Veterinaria, Pontificia Universidad Católica de Chile, Santiago, Chile

13

14

15

16

17

18 * **Correspondence:**
19 Corresponding Author
20 andrea.moreno@uc.cl

21

22

23 **Keywords: Antimicrobial resistance, Genomics, Clinically Relevant Bacteria, *E. coli*, *Klebsiella***
24 ***pneumoniae*, Food Safety.**

25 Abstract

26 Antimicrobial resistance is one of the most important public health concerns —it causes
 27 700,000 deaths annually according to the World Health Organization (WHO). Enterobacterales [such](#)
 28 [as](#) *E. coli* and *Klebsiella pneumoniae*, [have become](#) resistant to [many](#) relevant antimicrobials
 29 [including](#) carbapenems and extended spectrum cephalosporins. [These](#) clinically relevant resistant
 30 Enterobacterales (CRRE) [members are now](#) globally distributed in the environment including
 31 different food types (meats, produce, dairy). Unlike known foodborne pathogens, CRRE are not
 32 usually part of most food surveillance systems. However, numerous reports of CRRE highlight the
 33 importance of these bacteria in food and [have been shown to contribute to](#) the overall crisis of
 34 antimicrobial resistance. [This is especially important in the context of carriage of these pathogens by](#)
 35 [immuno-compromised individuals. CRRE infections upon consumption of contaminated food could](#)
 36 [colonize the human gastrointestinal tract and eventually be a source of systemic infections such as](#)
 37 [urinary tract infections or septicemia](#). While different aspects need to be considered to elucidate this,
 38 whole genome sequencing [along with metadata](#) could be used to understand genomic relationships of
 39 CRRE obtained from foods [and](#) humans, [including isolates from clinical](#) infections. Once robust
 40 scientific data is available on the role of CRRE in food, countries could move forward to better
 41 survey and control CRRE in food.

42 Introduction

43 Antimicrobial resistance (AMR) occurs when microorganisms evolve to resist the action of
 44 antimicrobials, making infections harder to treat, especially in critical patients leading to premature
 45 death (Dadgostar 2019). Unfortunately, bacteria have progressively acquired resistance to numerous
 46 antibiotics exhausting therapeutic alternatives and generating a serious public health problem and a
 47 worldwide alert (World Health Organization 2020). Importantly, AMR can jeopardize many of the
 48 advances of modern medicine, and threatens to turn usually treatable conditions into deadly infections
 49 that could carry a high death toll around the globe (Menichetti and Tagliaferri 2012; Chioro et al.,
 50 2015). Currently, AMR is estimated to cause 700,000 deaths per year, this number is projected to reach
 51 10 million deaths annually by 2050 (Hoffman et al., 2015).

52 The dynamics of AMR are complex and poorly understood. However, the connection between
 53 humans, animals, food, and the environment is increasingly recognized as a major player in the
 54 dissemination of AMR (Ludden et al., 2020). Therefore, the threat of AMR needs an integrated

[approach such as](#) the collaborative effort of key players from different disciplines at the WHO, Food and Agriculture Organization (FAO), and World Organization for Animal Health (OIE) tripartite (World Health Organization, 2021). To tackle AMR, it is important to reduce transmission of resistant bacteria [that are circulating among](#) humans and animals. [To accomplish this a better understanding of the role of these organisms in food production is needed. This knowledge will augment the responsible use of antibiotics in humans and animals](#) (Pokharel, Shrestha, and Adhikari 2020).

Food products are well-known carriers of foodborne antibiotic-resistant pathogens such as *Salmonella* spp. or *Campylobacter* spp. (Nordstrom et al., 2013; Allard et al., 2018). These organisms represent an important public health concern and are, therefore, subject to active surveillance in several countries (Hoffmann et al., 2017). Furthermore, food may also be contaminated with commensal organisms harboring antimicrobial resistance genes, most of these are not part of routine surveillance strategies (Hölzel et al., 2018). Importantly, several studies have already reported the presence of resistant bacteria in foods [but only a handful have established relationships between resistant bacteria in food and in humans at the same time.](#) This [risk is not comprehensively understood](#) much [less controlled](#) (Kim et al., 2015; Liu et al., 2018; Liu and Song 2019).

Clinically-Relevant Antimicrobial Resistant Bacteria

In 2017, the WHO published a priority list of antimicrobial resistant pathogens, highlighting the threat posed by multidrug-resistant bacteria (Harbarth et al., 2017). Common characteristics among these priority pathogens are their [wide](#) dissemination and their ability to horizontally acquire and transfer resistance traits (van Duin and Doi 2017; Logan and Weinstein 2017; Kelly et al., 2017). Importantly, all organisms regarded as ‘critical’ in WHO’s priority list are Gram-negatives and include carbapenem-resistant [bacteria of different taxa such as](#) *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, and Enterobacterales. The list also includes Enterobacterales that produce Extended Spectrum Beta-Lactamases (ESBL)—enzymes that can confer resistance to third-generation cephalosporins (Ghafourian et al., 2014).

Enterobacterales are particularly relevant because they can be found in numerous environments including food and cause significant human infections (Van Loon et al., 2018). For instance, infections caused by antibiotic-resistant *Klebsiella* spp. and *Escherichia coli* are associated with mortality rates of around 50% (Logan and Weinstein 2017; Lipari et al., 2020). The high

mortality caused by these pathogens is probably associated with the absence of effective treatments (Temkin et al., 2014). Globally, *Klebsiella pneumoniae* has been repeatedly identified over the past 30 years, as the most common Enterobacterales linked to the spread of ESBL genes in healthcare settings (Nordmann, Naas, and Poirel 2011). Importantly, patients infected with carbapenem-resistant *K. pneumoniae* have a 4-fold increased risk of death versus patients infected with antibiotic-susceptible *K. pneumoniae* (Ben-David et al., 2012). Therefore, carbapenem-resistant Enterobacterales along with those that produce ESBLs are considered clinically relevant-resistant bacteria, these are not only found in human but are also widely reported in the environment and food. However, the fate of clinically relevant resistant bacteria in food and their role in public health are not yet fully understood.

Food Reports and Subtyping of Clinically-Relevant Resistant Enterobacterales

Different food types have been found to contain clinically-relevant resistant Enterobacterales (CRRE)—specifically Enterobacterales producers of ESBL and/or carbapenemases (enzymes that confer resistance to carbapenems) (Liu et al., 2018; Zhang et al., 2019; Sugawara et al., 2019). Reports of CRRE in food are seen from many different countries; meats, produce, and dairy have all been found to harbor CRRE (Table 1). Most studies have focused on the study of CRRE in different meats such as chicken, beef, and pork; more incipient is the study of CRRE in produce and dairy (Table 1). Overall, the data suggest that CRRE appears to be widely and globally distributed in food.

E. coli and *K. pneumoniae* are the more commonly CRRE reported in food. These organisms are found to harbor genes coding for carbapenemases (e.g., *bla*_{OXA-48} and *bla*_{NDM}). These CRRE containing these genes have been found in meat and produce. CRRE carrying genes coding for the ESBL enzymes *bla*_{SHV}, *bla*_{CTX-M}, and *bla*_{TEM}, have been extensively reported in meat and produce (Table 1). To date, reports of CRRE in food have not necessarily facilitated our understanding of the public health role of the presence of CRRE in food and the risk for consumers of food-mediated acquisition of CRRE.

For decades the gold standard technique to understand the genetic relationships between bacteria has been pulsed-field gel electrophoresis (PFGE) (Salipante et al., 2015). Of the studies selected in Table 1, eight of them used PFGE to understand the relationships of CRRE isolates in food. These previous studies have shown the importance of subtyping isolates; the importance of obtaining greater resolution and summarizing genetic data can improved the study of CRRE in food.

For instance, one study used PFGE to type CRRE isolates obtained from vegetables and found two isolates co-carrying relevant resistant genes (i.e., the colistin resistant gene *mcr-1* and the β -lactamase *bla*NDM). These isolates were PFGE typed along with isolates of avian origin (Liu et al., 2017). However, the PFGE typing showed different patterns between avian and vegetable isolates, suggesting that vegetable isolates were not directly derived from the avian feces previously studied (Liu and Song 2019).

PFGE has a lower capacity to correctly identify strains in clusters than whole genome sequencing (WGS), which shows better resolution (Gona et al., 2020). To advance this knowledge, some studies have sequenced the genomes of CRRE obtained from food (Table 1); for instance, in Table 1 only 5 of the 28 studies included used genomic information. However, genomic relationships of CRRE isolated from food and CRRE obtained from human clinical cases or human colonization could improve our understanding of the public health risk of food-mediated acquisition of CRRE.

Whole Genome Sequencing as a tool to identify the public health risk of CRRE in food

Whole genome sequencing is a useful tool in food safety and public health facilitating our understanding of transmission and persistence of foodborne pathogens (Kovac et al., 2017). WGS has been crucial for studying foodborne outbreak investigations for almost a decade in the US and Europe (Brown et al., 2019; European Centre for Disease Prevention and Control, 2021). Numerous outbreaks have been rapidly resolved using WGS along with epidemiological data. For instance, in 2018 an outbreak of salmonellosis in the European Union related to eggs was resolved using WGS to confirm a source in Poland as the origin of the outbreak. This highlighted the public health value of international sharing of epidemiological data along with rapidly obtained WGS (Pijnacker et al., 2019). In addition, WGS have also been used to track contamination routes of *Listeria monocytogenes* in food production establishments (Nastasijevic et al., 2017). Therefore, WGS is now widely use in all sectors involved in food safety including governments, academia, and the food industry (Wang et al., 2016).

The use WGS is crucial to improve our understanding of the public health impact on CRRE in food. One study compared the genomes from *E. coli* obtained from humans, poultry, wild animals, and water collected in the past decade in Australia. Using WGS the study evaluated sequence types, mobile genetic elements, phylogroups, etc. to improve understanding of epidemiological clones and the presence of pervasive horizontal gene transfer across the species in all these sources (Touchon et

al., 2020). In this study, the authors compared around 1,300 strains to represent the genetic diversity of *E. coli* in Australia and looked for associations between the different phylogroups and the environment. Their results showed that the acquisition of mobile genetic elements (MGE) plays an important role in the first stages of genetic divergence between isolates, increasing the genome size of *E. coli*. The source of isolation would be determined by the number of MGEs acquired, reinforcing the idea that there is an association of certain genetic backgrounds with specific environments. Therefore, understanding the genetic context of each source of isolation is be a key step in understanding the movement of genes by including resistance and virulence genes at a specific location (Touchon et al., 2020). Consequently, it is very important to study the genomic relationships and MGEs of CRRE isolates found in food and to compare them with isolates obtained from human clinical cases or human colonization in healthy people.

To use WGS to infer genomic relationships of CRRE is not an easy task; mostly because the plausible effect of consumption of CRRE in food is not expected to represent an acute disease, but rather gastrointestinal colonization is proposed (Hölzel et al., 2018). Furthermore, the term foodborne urinary tract infection (FUTI) was also proposed as a new paradigm widening the perspective on the public health effects of CRRE in food (Nordstrom et al., 2013; Liu et al., 2018). WGS was used to study the genomic relationships of CRRE from food to humans using a One Health approach. A one-year study in Arizona compared the genomes of *E. coli* isolates from meat products and isolates from human urinary and blood sources (Liu et al., 2018). The study found one sublineage of *E. coli* ST131 found in poultry meat and human clinical cases raising the hypothesis of foodborne transmission and remarking the role of antimicrobial resistant *E. coli* in poultry meat (Liu et al., 2018). Importantly, the ability to cause UTI of *E. coli* isolates obtained from different meats was proven in murine models (Jakobsen et al., 2012).

Here, WGS has been an important tool to study the potential public health role of CRRE in food, and the whole transmission route is not completely understood. The variety of food types in which CRRE have been reported include food that is not further processed at home (e.g., fruits and vegetables). To better understand this, CRRE obtained from different food types need to be studied along with human gastrointestinal colonization and clinical cases (Fig. 1). WGS can provide information about virulence potential, adaptation and survival such as resistance to biocides, metals, and antimicrobial drugs along with the plasticity of genomes of food isolates. For example, in antibiotic-resistant *Salmonella* obtained from retail meats and clinical isolates, WGS identified the

[resistance genes and the genomic environment of these genes](#) (Keefer et al., 2018) [providing unique and valuable information for the study of the spread of antimicrobial resistance. Similarly, virulence-associated genes of ESBL-producing *K. pneumoniae* from different sources including vegetables has been identification by using WGS](#) (Chi et al., 2019). [Both examples provide an enormous advance and facilitate the detection of potential risks to public health.](#)

In Chile, the Millennium Initiative for Collaborative Research on Bacterial Resistance (microb-r.org; which is a multi-institution research center focusing on the study of AMR) partnered with the Maule Cohort (MAUCO) (Ferreccio et al., 2016; 2020). Here, suitable conditions are given to [study AMR and to](#) use WGS of CRRE obtained from different food types and from human volunteers (10,000 adults inhabitants) from [Molina county](#) where the MAUCO cohort is located. MAUCO cohort's participants represent a unique opportunity to advance the current understanding on the public health consequences of the presence of CRRE in the food supply. Close genomic relatedness of CRRE from different sources could help to trace isolates from the food supply to human gastrointestinal tracts, to understand the movements of mobile elements as plasmids and resistance genes in CRRE from different food and water sources, and to infer phylogenetic relationships ([Fig. 1](#)). Colonization is one aspect but [clones](#) that are capable of causing bloodstream infections can also be traced to gastrointestinal colonization and to food using WGS. Identification of clonal isolates in food, in colonized participants, and finally in bloodstream infections could possible close the current gap and provide important data to policy makers.

[The isolation of CRRE from samples from three main compartments are necessary](#) to obtain relevant data ([Fig. 1](#)). [Importantly, WGS of isolated CRRE along with metadata from the cohort participants obtained through, for instance questionnaires or participants records, could provide information on susceptible populations related to foodborne acquisition of CRRE. The elderly or immuno-compromised individuals are more susceptible to foodborne infections](#) (Mohácsi-Farkas 2016), and thus we expect that [this is particularly important to them. Moreover, this approach can also include the impact of chronic diseases \(e.g., diabetes, hypertension\), along with risk factors such as antimicrobial therapy, patterns of food consumption, among others](#) ([Fig. 1](#)).

Discussion

The clinical interest in antimicrobial-resistant bacteria strikes at the hospitals where therapeutic options to treat infections caused by, for instance carbapenem resistant or ESBL producer

Enterobacterias are limited (Rodriguez-Baño et al., 2018). Reports on the detection of bacteria resistant to clinically relevant antibiotics [associated with](#) food have increased worldwide, however, the outcome of CRRE in different food types and the potential public health impacts are currently unknown. Colonization with carbapenem resistant or ESBL-producer [Enterobacterales](#) has been described as one plausible outcome (Hölzel et al., 2018). The consumption of food harboring CRRE could further colonize the human gut—this will not necessarily cause or either be associated with a known foodborne disease. [In addition, colonization could further worsen some conditions](#), e.g., [recent evidence has shown that individuals colonized by multidrug resistant microorganisms are more likely to present systemic infections such as septicemia with a higher chance of treatment failure than not colonized](#) ones (Ferreira et al., 2018). Several questions remain and important knowledge gaps need to be filled. These include a better understanding on the dose-response (Hölzel et al., 2018) and a genomic-based approach that better explains the distribution and closeness with resistant bacteria present in the human gastrointestinal tract with bacteria obtained from different food types.

[Many](#) antibiotics of human interest are [used](#) in domestic animals and may reach food and the environment, although AMR in food and the role in the overall AMR crisis are difficult to compare due to a lack of [published studies or surveillance programs](#) in animal, environmental, [or food](#) samples. [For instance only a small fraction of food types have been studied as carriers of CRRE](#) (Schrijver et al., 2017; Hudson et al., 2017). To avoid this problem, it is recommended to design studies with statistical support to determine the number of samples and genomes needed. It is also necessary to use methodologies that make relationships between what has been found in animals, food, and the environment with all the antecedents that currently exist in humans (the most common studied). Here, the use of study cohorts is recommended and can help to explain risk factors, the origin of some resistance plasmids or multidrug resistant bacteria, CRRE, etc. (Arcilla et al. 2017; Barría 2018).

Genomic tools and longitudinal study design at a global level, can improve food safety and ensure a good surveillance system that is reliable, reproducible, and sustainable. This promotes collaborative studies among different regions or countries. Importantly, the developed countries could work together with underdeveloped countries, which do not have the financing or tools necessary to carry out this type of study and which are often suppliers of food of animal or vegetable origin (Errecart 2015; ONU, 2019).

Finally, we believe that, in addition to the development of new antimicrobials, it is of extreme importance to carry out the follow-up and analysis of carbapenem-resistant and ESBL-producing Enterobacterales. The WGS, the generation of worldwide databases, and the publication of standard protocols for working with these pathogens in food are needed.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author Contributions

CDG, FPA, AIMS wrote the manuscript. JMM, SC critically reviewed the manuscript.

Funding

This work was funded by FONDECYT 1181167 and by the ANID Millennium Science Initiative/ Millennium Initiative for Collaborative Research on Bacterial Resistance, MICROB-R, NCN17_081.

Acknowledgments

Figure 1 was created with BioRender.com

References

- Abdallah, H. M., E. A. Reuland, B. B. Wintermans, N. Al Naiemi, A. Koek, A. M. Abdelwahab, A. M. Ammar, A. A. Mohamed, and C. M.J.E. Vandenbroucke-Grauls. 2015. "Extended-Spectrum β -Lactamases and/or Carbapenemases-Producing *Enterobacteriaceae* Isolated from Retail Chicken Meat in Zagazig, Egypt." *PLoS ONE* 10 (8): 1–8. <https://doi.org/10.1371/journal.pone.0136052>.
- Allard, Marc W., Rebecca Bell, Christina M. Ferreira, Narjol Gonzalez-Escalona, Maria Hoffmann, Tim Muruvanda, Andrea Ottesen, et al. 2018. "Genomics of Foodborne Pathogens for Microbial Food Safety." *Current Opinion in Biotechnology*. <https://doi.org/10.1016/j.copbio.2017.11.002>.
- Arcilla, Maris S., Jarne M. van Hattem, Manon R. Haverkate, Martin C.J. Bootsma, Perry J.J. van Genderen, Abraham Goorhuis, Martin P. Grobusch, et al. 2017. "Import and Spread of Extended-Spectrum β -Lactamase-Producing *Enterobacteriaceae* by International Travellers

- (COMBAT Study): A Prospective, Multicentre Cohort Study.” *The Lancet Infectious Diseases* 17 (1): 78–85. [https://doi.org/10.1016/S1473-3099\(16\)30319-X](https://doi.org/10.1016/S1473-3099(16)30319-X).
- Barria, René Mauricio. 2018. “Introductory Chapter: The Contribution of Cohort Studies to Health Sciences.” In *Cohort Studies in Health Sciences*. <https://doi.org/10.5772/intechopen.80178>.
- Ben-David, D., R. Kordevani, N. Keller, I. Tal, A. Marzel, O. Gal-Mor, Y. Maor, and G. Rahav. 2012. “Outcome of Carbapenem Resistant *Klebsiella pneumoniae* Bloodstream Infections.” *Clinical Microbiology and Infection*. <https://doi.org/10.1111/j.1469-0691.2011.03478.x>.
- Bhutani, Natasha, Chithra Muraleedharan, Deepa Talreja, Sonia Walia Rana, Sandeep Walia, Ashok Kumar, and Satish K. Walia. 2015. “Occurrence of Multidrug Resistant Extended Spectrum Beta-Lactamase-Producing Bacteria on Iceberg Lettuce Retailed for Human Consumption.” *BioMed Research International*. <https://doi.org/10.1155/2015/547547>.
- Botelho, Larissa Alvarenga Batista, Gabriela Bergiante Kraychete, Patrícia Batista Rocha, Ana Paula De Souza Da-Silva, Renata Cristina Picão, Beatriz Meurer Moreira, and Raquel Regina Bonelli. 2020. “CTX-M- And PAMP-C-Encoding Genes Are Associated with Similar Mobile Genetic Elements in *Escherichia coli* Isolated from Different Brands of Brazilian Chicken Meat.” *Microbial Drug Resistance*. <https://doi.org/10.1089/mdr.2019.0043>.
- Brown, Eric, Uday Dessai, Sherri McGarry, and Peter Gerner-smidt. 2019. “Use of Whole-Genome Sequencing for Food Safety and Public Health in the United States” 16 (7): 441–50. <https://doi.org/10.1089/fpd.2019.2662>.
- Chi, Xiaohui, Björn Berglund, Huiyun Zou, Beiwen Zheng, Stefan Börjesson, Xiang Ji, Jakob Ottoson, Cecilia Stålsby Lundborg, Xuewen Li, and Lennart E. Nilsson. 2019. “Strains of Extended-Spectrum β -Lactamase-Producing *Klebsiella pneumoniae* Occurring in Environmental Sources in a Rural Area of China by Using Whole-Genome Sequencing.” *Frontiers in Microbiology*. <https://doi.org/10.3389/fmicb.2019.00211>.
- Chioro, Arthur, Awa Marie Coll-Seck, Bent Høie, Nila Moeloek, Aaron Motsoaledi, Rajata Rajatanavin, and Marisol Touraine. 2015. “Antimicrobial Resistance: A Priority for Global Health Action.” *Bulletin of the World Health Organization* 93 (7): 439. <https://doi.org/10.2471/BLT.15.158998>.
- Clemente, Lurdes, Vera Manageiro, Ivone Correia, Ana Amaro, Teresa Albuquerque, Patrícia

- Themudo, Eugénia Ferreira, and Manuela Caniça. 2019. “Revealing Mcr-1-Positive ESBL-Producing *Escherichia Coli* Strains among *Enterobacteriaceae* from Food-Producing Animals (Bovine, Swine and Poultry) and Meat (Bovine and Swine), Portugal, 2010–2015.” *International Journal of Food Microbiology*. <https://doi.org/10.1016/j.ijfoodmicro.2019.02.006>.
- Dadgostar, Porooshat. 2019. “Antimicrobial Resistance: Implications and Costs.” *Infection and Drug Resistance* 12: 3903–10. <https://doi.org/10.2147/IDR.S234610>.
- Duin, David van, and Yohei Doi. 2017. “The Global Epidemiology of Carbapenemase-Producing *Enterobacteriaceae*.” *Virulence*. <https://doi.org/10.1080/21505594.2016.1222343>.
- Errecart, Valeria. 2015. “Análisis Del Mercado Mundial de Carnes.” *Cere* 1: 35. http://www.unsam.edu.ar/escuelas/economia/economia_regional/CERE - Mayo - 2015.pdf.
- European Centre for Disease Prevention and Control (ECDC). 2021. *ECDC Strategic Framework for the Integration of Molecular and Genomic Typing into European Surveillance and Multi-Country Outbreak Investigations*.
- Ferreccio, Catterina, Andrea Huidobro, Sandra Cortés, Claudia Bambs, Pablo Toro, Vanessa Van De Wyngard, Johanna Acevedo, et al. 2020. “Cohort Profile: The Maule Cohort (MAUCO).” *International Journal of Epidemiology*. <https://doi.org/10.1093/ije/dyaa003>.
- Ferreccio, Catterina, Juan Carlos Roa, Claudia Bambs, Alejandra Vives, Alejandro H. Corvalán, Sandra Cortés, Claudia Foerster, et al. 2016. “Study Protocol for the Maule Cohort (MAUCO) of Chronic Diseases, Chile 2014-2024.” *BMC Public Health*. <https://doi.org/10.1186/s12889-015-2454-2>.
- Ferreira, A. M., F. Moreira, T. Guimaraes, F. Spadão, J. F. Ramos, M. V. Batista, J. S. Filho, S. F. Costa, and V. Rocha. 2018. “Epidemiology, Risk Factors and Outcomes of Multi-Drug-Resistant Bloodstream Infections in Haematopoietic Stem Cell Transplant Recipients: Importance of Previous Gut Colonization.” *Journal of Hospital Infection*. <https://doi.org/10.1016/j.jhin.2018.03.004>.
- Garcia-Graells, Cristina, Bas Berbers, Bavo Verhaegen, Kevin Vanneste, Kathleen Marchal, Nancy H.C. Roosens, Nadine Botteldoorn, and Sigrid C.J. De Keersmaecker. 2020. “First Detection of a Plasmid Located Carbapenem Resistant BlaVIM-1 Gene in *E. coli* Isolated from Meat Products at Retail in Belgium in 2015.” *International Journal of Food Microbiology*.

- 322 <https://doi.org/10.1016/j.ijfoodmicro.2020.108624>.
- 323 Ghafourian, Sobhan, Nourkhoda Sadeghifard, Sara Soheili, and Zamberi Sekawi. 2014. "Extended
324 Spectrum Beta-Lactamases: Definition, Classification and Epidemiology." *Current Issues in*
325 *Molecular Biology*. <https://doi.org/10.21775/cimb.017.011>.
- 326 Gona, Floriana, Francesco Comandatore, Simone Battaglia, Aurora Piazza, Alberto Trovato,
327 Giovanni Lorenzin, Paola Cichero, et al. 2020. "Comparison of Core-Genome MLST, CoreSNP
328 and PFGE Methods for *Klebsiella pneumoniae* Cluster Analysis." *Microbial Genomics*.
329 <https://doi.org/10.1099/mgen.0.000347>.
- 330 Harbarth, S, G Kahlmeter, J Kluytmans, M Mendelson, Groote Schuur Hospital, Cape Town, South
331 Africa, et al. 2017. "Global priority list of antibiotic-resistant bacteria to guide research,
332 discovery, and development of new antibiotics"
- 333 Hassen, Bilel, Mohamed Salah Abbassi, Laura Ruiz-Ripa, Olouwafemi M. Mama, Abdennaceur
334 Hassen, Carmen Torres, and Salah Hammami. 2020. "High Prevalence of Mcr-1 Encoding
335 Colistin Resistance and First Identification of BlaCTX-M-55 in ESBL/CMY-2-Producing
336 *Escherichia coli* Isolated from Chicken Faeces and Retail Meat in Tunisia." *International*
337 *Journal of Food Microbiology* 318 (October 2019): 108478.
338 <https://doi.org/10.1016/j.ijfoodmicro.2019.108478>.
- 339 Hayashi, Wataru, Yusuke Ohsaki, Yui Taniguchi, Shota Koide, Kumiko Kawamura, Masahiro
340 Suzuki, Kouji Kimura, et al. 2018. "High Prevalence of BlaCTX-M-14 among Genetically
341 Diverse *Escherichia coli* Recovered from Retail Raw Chicken Meat Portions in Japan."
342 *International Journal of Food Microbiology*. <https://doi.org/10.1016/j.ijfoodmicro.2018.08.003>.
- 343 Hoang, T. A.V., T. N.H. Nguyen, S. Ueda, Q. P. Le, T. T.N. Tran, T. N.D. Nguyen, T. V.K. Dao, et
344 al. 2017. "Common Findings of Bla CTX-M-55-Encoding 104–139 Kbp Plasmids Harbored by
345 Extended-Spectrum β -Lactamase-Producing *Escherichia coli* in Pork Meat, Wholesale Market
346 Workers, and Patients with Urinary Tract Infection in Vietnam." *Current Microbiology*.
347 <https://doi.org/10.1007/s00284-016-1174-x>.
- 348 Hoffman, Steven J., Kevin Outterson, John Arne Røttingen, Otto Cars, Charles Clift, Zain Rizvi,
349 Fiona Rotberg, Göran Tomson, and Anna Zorzet. 2015. "An International Legal Framework to
350 Address Antimicrobial Resistance." *Bulletin of the World Health Organization* 93 (2): 66.

- 351 <https://doi.org/10.2471/BLT.15.152710>.
- 352 Hoffmann, Sandra, Brecht Devleesschauwer, Willy Aspinall, Roger Cooke, Tim Corrigan, Arie
 353 Havelaar, Frederick Angulo, et al. 2017. "Attribution of Global Foodborne Disease to Specific
 354 Foods: Findings from a World Health Organization Structured Expert Elicitation." *PLoS ONE*.
 355 <https://doi.org/10.1371/journal.pone.0183641>.
- 356 Hölzel, Christina Susanne, Julia Louisa Tetens, and Karin Schwaiger. 2018. "Unraveling the Role of
 357 Vegetables in Spreading Antimicrobial-Resistant Bacteria: A Need for Quantitative Risk
 358 Assessment." *Foodborne Pathogens and Disease*. <https://doi.org/10.1089/fpd.2018.2501>.
- 359 Hudson, John A., Lynn J. Frewer, Glyn Jones, Paul A. Brereton, Mark J. Whittingham, and Gavin
 360 Stewart. 2017. "The Agri-Food Chain and Antimicrobial Resistance: A Review." *Trends in*
 361 *Food Science and Technology*. <https://doi.org/10.1016/j.tifs.2017.09.007>.
- 362 Huizinga, Pepijn, Marjolein Kluytmans Van Den Bergh, John W. Rossen, Ina Willemsen, Carlo
 363 Verhulst, Paul H.M. Savelkoul, Alexander W. Friedrich, Silvia García-Cobos, and Jan
 364 Kluytmans. 2019. "Decreasing Prevalence of Contamination with Extended-Spectrum Beta-
 365 Lactamaseproducing *Enterobacteriaceae* (ESBL-E) in Retail Chicken Meat in the Netherlands." *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0226828>.
- 367 Jakobsen, L., P. Garneau, G. Bruant, J. Harel, S. S. Olsen, L. J. Porsbo, A. M. Hammerum, and N.
 368 Frimodt-Møller. 2012. "Is *Escherichia coli* Urinary Tract Infection a Zoonosis? Proof of Direct
 369 Link with Production Animals and Meat." *European Journal of Clinical Microbiology and*
 370 *Infectious Diseases*. <https://doi.org/10.1007/s10096-011-1417-5>.
- 371 Kaesbohrer, Annemarie, Karin Bakran-Lebl, Alexandra Irrgang, Jennie Fischer, Peter Kämpf, Arthur
 372 Schiffmann, Christiane Werckenthin, Matthias Busch, Lothar Kreienbrock, and Katja Hille.
 373 2019. "Diversity in Prevalence and Characteristics of ESBL/PAmpC Producing *E. coli* in Food
 374 in Germany." *Veterinary Microbiology* 233 (January 2018): 52–60.
 375 <https://doi.org/10.1016/j.vetmic.2019.03.025>.
- 376 Keefer, Andrea B., Lingzi Xiaoli, Nkuchia M. M'ikanatha, Kuan Yao, Maria Hoffmann, and Edward
 377 G. Dudley. 2018. "Retrospective Whole-Genome Sequencing Analysis Distinguished PFGE and
 378 Drug Resistance Matched Retail Meat and Clinical Salmonella Isolates." *BioRxiv*.
 379 <https://doi.org/10.1101/356857>.

- 380 Kelly, Ana M., Barun Mathema, and Elaine L. Larson. 2017. "Carbapenem-Resistant
381 *Enterobacteriaceae* in the Community: A Scoping Review." *International Journal of*
382 *Antimicrobial Agents*. <https://doi.org/10.1016/j.ijantimicag.2017.03.012>.
- 383 Kim, Hong Seok, Jung Whan Chon, Young Ji Kim, Dong Hyeon Kim, Mu sang Kim, and Kun Ho
384 Seo. 2015. "Prevalence and Characterization of Extended-Spectrum- β -Lactamase-Producing
385 *Escherichia coli* and *Klebsiella pneumoniae* in Ready-to-Eat Vegetables." *International Journal*
386 *of Food Microbiology*. <https://doi.org/10.1016/j.ijfoodmicro.2015.04.049>.
- 387 Kluytmans-van den bergh, Marjolein, P. Huizinga, M. J. Bonten, M. Bos, K. De Bruyne, A. W.
388 Friedrich, J. W. Rossen, P. H. Savelkoul, and J. A. Kluytmans. 2016. "Presence of Mcr-1-
389 Positive *Enterobacteriaceae* in Retail Chicken Meat but Not in Humans in the Netherlands since
390 2009." *Eurosurveillance*. <https://doi.org/10.2807/1560-7917.ES.2016.21.9.30149>.
- 391 Kovac, Jasna, Henk Den Bakker, Laura M Carroll, and Martin Wiedmann. 2017. "Trends in
392 Analytical Chemistry Precision Food Safety : A Systems Approach to Food Safety Facilitated
393 by Genomics Tools." *Trends in Analytical Chemistry* 96: 52–61.
394 <https://doi.org/10.1016/j.trac.2017.06.001>.
- 395 Li, Lili, Lei Ye, Li Yu, Chenqing Zhou, and Hecheng Meng. 2016. "Characterization of Extended
396 Spectrum B-Lactamase Producing Enterobacteria and Methicillin-Resistant *Staphylococcus*
397 *aureus* Isolated from Raw Pork and Cooked Pork Products in South China." *Journal of Food*
398 *Science*. <https://doi.org/10.1111/1750-3841.13346>.
- 399 Lipari, Flavio G., Daniela Hernández, Mario Vilaró, Juan P. Caeiro, and Héctor Alex Saka. 2020.
400 "Caracterización Clínica, Epidemiológica y Microbiológica de Bacteriemias Producidas Por
401 Enterobacterias Resistentes a Carbapenems En Un Hospital Universitario de Córdoba,
402 Argentina." *Revista Chilena de Infectología*. [https://doi.org/10.4067/s0716-](https://doi.org/10.4067/s0716-10182020000400362)
403 [10182020000400362](https://doi.org/10.4067/s0716-10182020000400362).
- 404 Liu, Bao Tao, and Feng Jing Song. 2019. "Emergence of Two *Escherichia coli* Strains Co-Harboring
405 Mcr-1 and BlaNDM in Fresh Vegetables from China." *Infection and Drug Resistance*.
406 <https://doi.org/10.2147/IDR.S211746>.
- 407 Liu, Bao Tao, Feng Jing Song, Ming Zou, Qi Di Zhang, and Hu Shan. 2017. "High Incidence of
408 *Escherichia coli* Strains Coharboring Mcr-1 and BlaNDM from Chickens." *Antimicrobial*

- 409 *Agents and Chemotherapy*. <https://doi.org/10.1128/AAC.02347-16>.
- 410 Liu, Bao Tao, Xiao Yan Zhang, Shu Wei Wan, Jun Jie Hao, Rui De Jiang, and Feng Jing Song. 2018.
411 “Characteristics of Carbapenem-Resistant *Enterobacteriaceae* in Ready-to-Eat Vegetables in
412 China.” *Frontiers in Microbiology*. <https://doi.org/10.3389/fmicb.2018.01147>.
- 413 Liu, Cindy M, Marc Stegger, Maliha Aziz, Timothy J Johnson, Kara Waits, Lora Nordstrom, Lori
414 Gauld, et al. 2018. “*Escherichia coli* ST131-H22 as a Foodborne Uropathogen.” *MBio*.
- 415 Logan, Latania K., and Robert A. Weinstein. 2017. “The Epidemiology of Carbapenem-Resistant
416 *Enterobacteriaceae*: The Impact and Evolution of a Global Menace.” *Journal of Infectious*
417 *Diseases*. <https://doi.org/10.1093/infdis/jiw282>.
- 418 Loon, Karlijn Van, Anne F. Voor In’T Holt, and Margreet C. Vos. 2018. “A Systematic Review and
419 Meta-Analyses of the Clinical Epidemiology of Carbapenem-Resistant *Enterobacteriaceae*.”
420 *Antimicrobial Agents and Chemotherapy*. <https://doi.org/10.1128/AAC.01730-17>.
- 421 Ludden, Catherine, Danesh Moradigaravand, Dorota Jamroz, Theodore Gouliouris, Beth Blane,
422 Plamena Naydenova, Juan Hernandez-Garcia, et al. 2020. “A One Health Study of the Genetic
423 Relatedness of *Klebsiella pneumoniae* and Their Mobile Elements in the East of England.”
424 *Clinical Infectious Diseases*. <https://doi.org/10.1093/cid/ciz174>.
- 425 Menichetti, Francesco, and Enrico Tagliaferri. 2012. “Antimicrobial Resistance in Internal Medicine
426 Wards.” *Internal and Emergency Medicine* 7 (SUPPL. 3): 271–81.
427 <https://doi.org/10.1007/s11739-012-0828-3>.
- 428 Mohácsi-Farkas, Csilla. 2016. “Food Irradiation: Special Solutions for the Immuno-Compromised.”
429 *Radiation Physics and Chemistry*. <https://doi.org/10.1016/j.radphyschem.2016.06.033>.
- 430 Montso, Kotsoana Peter, Sicelo Beauty Dlamini, Ajay Kumar, Collins Njie Ateba, and Herney A.
431 Garcia-Perdomo. 2019. “Antimicrobial Resistance Factors of Extended-Spectrum Beta-
432 Lactamases Producing *Escherichia coli* and *Klebsiella pneumoniae* Isolated from Cattle Farms
433 and Raw Beef in North-West Province, South Africa.” *BioMed Research International*.
434 <https://doi.org/10.1155/2019/4318306>.
- 435 Nastasijevic, Ivan, Dubravka Milanov, Branko Velebit, Vesna Djordjevic, Craig Swift, Anaïs
436 Painset, and Brankica Lakicevic. 2017. “International Journal of Food Microbiology Tracking of

- 437 *Listeria monocytogenes* in Meat Establishment Using Whole Genome Sequencing as a Food
 438 Safety Management Tool : A Proof of Concept.” *International Journal of Food Microbiology*
 439 257 (March): 157–64. <https://doi.org/10.1016/j.ijfoodmicro.2017.06.015>.
- 440 Nordmann, Patrice, Thierry Naas, and Laurent Poirel. 2011. “Global Spread of Carbapenemase
 441 Producing *Enterobacteriaceae*.” *Emerging Infectious Diseases*.
 442 <https://doi.org/10.3201/eid1710.110655>.
- 443 Nordstrom, Lora, Cindy M. Liu, and Lance B. Price. 2013. “Foodborne Urinary Tract Infections: A
 444 New Paradigm for Antimicrobial-Resistant Foodborne Illness.” *Frontiers in Microbiology*.
 445 <https://doi.org/10.3389/fmicb.2013.00029>.
- 446 Organización Naciones Unidas. 2019. “Récord de Economías Dependientes de La Exportación de
 447 Productos Básicos.” Asuntos Económicos. 2019. <https://news.un.org/es/story/2019/05/1455981>.
- 448 Pehlivanlar Önen, Sevda, Özkan Aslantaş, Ebru Şebnem Yılmaz, and Cemil Kürekci. 2015.
 449 “Prevalence of β -Lactamase Producing *Escherichia coli* from Retail Meat in Turkey.” *Journal*
 450 *of Food Science*. <https://doi.org/10.1111/1750-3841.12984>.
- 451 Pijnacker, Roan, Timothy J Dallman, Aloys S L Tijsma, Gillian Hawkins, Lesley Larkin, Saara M
 452 Kotila, Giusi Amore, and Ettore Amato. 2019. “Articles An International Outbreak of
 453 *Salmonella enterica* Serotype Enteritidis Linked to Eggs from Poland : A Microbiological and
 454 Epidemiological Study” 3099 (19): 1–9. [https://doi.org/10.1016/S1473-3099\(19\)30047-7](https://doi.org/10.1016/S1473-3099(19)30047-7).
- 455 Pokharel, Sunil, Priyanka Shrestha, and Bipin Adhikari. 2020. “Antimicrobial Use in Food Animals
 456 and Human Health: Time to Implement ‘One Health’ Approach.” *Antimicrobial Resistance and*
 457 *Infection Control* 9 (1): 1–5. <https://doi.org/10.1186/s13756-020-00847-x>.
- 458 Randall, L. P., M. P. Lodge, N. C. Elviss, F. L. Lemma, K. L. Hopkins, C. J. Teale, and N.
 459 Woodford. 2017. “Evaluation of Meat, Fruit and Vegetables from Retail Stores in Five United
 460 Kingdom Regions as Sources of Extended-Spectrum Beta-Lactamase (ESBL)-Producing and
 461 Carbapenem-Resistant *Escherichia coli*.” *International Journal of Food Microbiology* 241:
 462 283–90. <https://doi.org/10.1016/j.ijfoodmicro.2016.10.036>.
- 463 Rasmussen, Mette Marie, Japheth A. Opintan, Niels Frimodt-Møller, and Bjarne Styris have. 2015.
 464 “Beta-Lactamase Producing *Escherichia coli* Isolates in Imported and Locally Produced
 465 Chicken Meat from Ghana.” *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0139706>.

- Rebbah, Nesrine, Yamina Messai, Pierre Châtre, Marisa Haenni, Jean Yves Madec, and Rabah Bakour. 2018. "Diversity of CTX-M Extended-Spectrum β -Lactamases in *Escherichia coli* Isolates from Retail Raw Ground Beef: First Report of CTX-M-24 and CTX-M-32 in Algeria." *Microbial Drug Resistance*. <https://doi.org/10.1089/mdr.2017.0171>.
- Reuland, E. A., N. al Naiemi, S. A. Raadsen, P. H.M. Savelkoul, J. A.J.W. Kluytmans, and C. M.J.E. Vandenbroucke-Grauls. 2014. "Prevalence of ESBL-Producing *Enterobacteriaceae* in Raw Vegetables." *European Journal of Clinical Microbiology and Infectious Diseases*. <https://doi.org/10.1007/s10096-014-2142-7>.
- Rodriguez-Baño, Jesús, Belen Gutiérrez-Gutiérrez, Isabel Machuca, and Alvaro Pascual. 2018. "Treatment of Infections Caused by Extended-Spectrum-Beta-Lactamase-, AmpC-, and Carbapenemase-Producing *Enterobacteriaceae*." *Clinical Microbiology Reviews*. DOI:10.1128/CMR.00079-17
- Saidani, Mariem, Lilia Messadi, Alya Soudani, Monia Daaloul-Jedidi, Pierre Châtre, Faten Ben Chehida, Aymen Mamlouk, Wassim Mahjoub, Jean Yves Madec, and Marisa Haenni. 2018. "Epidemiology, Antimicrobial Resistance, and Extended-Spectrum Beta-Lactamase-Producing *Enterobacteriaceae* in Clinical Bovine Mastitis in Tunisia." *Microbial Drug Resistance*. <https://doi.org/10.1089/mdr.2018.0049>.
- Salipante, Stephen J., Dhruva J. SenGupta, Lisa A. Cummings, Tyler A. Land, Daniel R. Hoogestraat, and Brad T. Cookson. 2015. "Application of Whole-Genome Sequencing for Bacterial Strain Typing in Molecular Epidemiology." *Journal of Clinical Microbiology*. <https://doi.org/10.1128/JCM.03385-14>.
- Schill, Franziska, Amir Abdulmawjood, Günter Klein, and Felix Reich. 2017. "Prevalence and Characterization of Extended-Spectrum β -Lactamase (ESBL) and AmpC β -Lactamase Producing *Enterobacteriaceae* in Fresh Pork Meat at Processing Level in Germany." *International Journal of Food Microbiology*. <https://doi.org/10.1016/j.ijfoodmicro.2017.06.010>.
- Schrijver, R., M. Stijntjes, J. Rodríguez-Baño, E. Tacconelli, N. Babu Rajendran, and A. Voss. 2017. "Review of Antimicrobial Resistance Surveillance Programmes in Livestock and Their Meat in Europe, with a Focus on Antimicrobial Resistance Patterns in Humans." *Clinical Microbiology and Infection*. <https://doi.org/10.1016/j.cmi.2017.09.013>.

- 495 Sugawara, Yo, Hideharu Hagiya, Yukihiro Akeda, Mya Mya Aye, Hpoo Pwint Myo Win, Noriko
496 Sakamoto, Rathina Kumar Shanmugakani, et al. 2019. "Dissemination of Carbapenemase-
497 Producing *Enterobacteriaceae* Harboursing Bla NDM or Bla IMI in Local Market Foods of
498 Yangon, Myanmar." *Scientific Reports*. <https://doi.org/10.1038/s41598-019-51002-5>.
- 499 Tadesse, Daniel A., Cong Li, Sampa Mukherjee, Chih Hao Hsu, Sonya Bodeis Jones, Stuart A.
500 Gaines, Claudine Kabera, et al. 2018. "Whole-Genome Sequence Analysis of CTX-M
501 Containing *Escherichia coli* Isolates from Retail Meats and Cattle in the United States."
502 *Microbial Drug Resistance*. <https://doi.org/10.1089/mdr.2018.0206>.
- 503 Tekiner, İsmail Hakkı, and Haydar Özpınar. 2016. "Occurrence and Characteristics of Extended
504 Spectrum Beta-Lactamases-Producing *Enterobacteriaceae* from Foods of Animal Origin."
505 *Brazilian Journal of Microbiology*. <https://doi.org/10.1016/j.bjm.2015.11.034>.
- 506 Temkin, Elizabeth, Amos Adler, Anat Lerner, and Yehuda Carmeli. 2014. "Carbapenem-Resistant
507 *Enterobacteriaceae*: Biology, Epidemiology, and Management." *Annals of the New York
508 Academy of Sciences*. <https://doi.org/10.1111/nyas.12537>.
- 509 Touchon, Marie, Amandine Perrin, Jorge Andre Moura De Sousa, Belinda Vangchhia, Samantha
510 Burn, Claire L. O'Brien, Erick Denamur, David Gordon, and Eduardo P.C. Rocha. 2020.
511 Phylogenetic Background and Habitat Drive the Genetic Diversification of *Escherichia coli*.
512 *PLoS Genetics*. Vol. 16. <https://doi.org/10.1371/journal.pgen.1008866>.
- 513 Wang, Siyun, Daniel Weller, Justin Falardeau, Laura K Strawn, O Fernando, Aiko D Adell, and
514 Andrea I Moreno Switt. 2016. "Food Safety Trends: From Globalization of Whole Genome
515 Sequencing to Application of New Tools to Prevent Foodborne Diseases." *Trends in Food
516 Science & Technology*. <https://doi.org/10.1016/j.tifs.2016.09.016>.
- 517 World Health Organization. n.d. "FAO/OIE/WHO Tripartite Collaboration on AMR." Food Safety.
518 Accessed February 4, 2021. [https://www.who.int/foodsafety/areas_work/antimicrobial-
519 resistance/tripartite/en/](https://www.who.int/foodsafety/areas_work/antimicrobial-resistance/tripartite/en/).
- 520 World Health Organization (WHO). 2020. "Antimicrobial Resistance." 2020.
521 <https://www.who.int/news-room/fact-sheets/detail/antimicrobial-resistance>.
- 522 Ye, Qinghua, Qingping Wu, Shuhong Zhang, Jumei Zhang, Guangzhu Yang, Juan Wang, Liang Xue,
523 and Moutong Chen. 2018. "Characterization of Extended-Spectrum β -Lactamase-Producing

- 524 *Enterobacteriaceae* from Retail Food in China.” *Frontiers in Microbiology*.
 525 <https://doi.org/10.3389/fmicb.2018.01709>.
- 526 Zhang, Qianhui, Luchao Lv, Xiuyu Huang, Ying Huang, Zilin Zhuang, Jiaxun Lu, Enyu Liu, et al.
 527 2019. “Rapid Increase in Carbapenemase-Producing *Enterobacteriaceae* in Retail Meat Driven
 528 by the Spread of the BlaNDM-5-Carrying IncX3 Plasmid in China from 2016 to 2018.”
 529 *Antimicrobial Agents and Chemotherapy*. <https://doi.org/10.1128/AAC.00573-19>.
- 530

In review

531 **Figure 1. Schematic representation of use of whole genome sequencing to determine genetic**
532 **relatedness of clinically relevant resistant Enterobacterales (CRRE) presence in food and**
533 **humans. A) Samples from three distinct origins including different food types, human samples of**
534 **cohort participants colonized in the community and samples from hospitalized patients with a**
535 **bloodstream infection. B) Isolation of CRRE from the origins in A along with data collection. C)**
536 **Expected results include identification of mobile elements and phylogenetic relatedness on CRRE**
537 **from these three origins, and prediction of risk factors using computational analysis.**

In review

538 **Table 1.** Selected reports of Clinically relevant antimicrobial resistant bacteria in different foods.

Bacterial Species reported	Food type	Carbapenems, cephalosporins and colistin resistance genes reported ^{2,3}	PFGE/Genome Sequenced	Country	Reference
Animal based food					
<i>E. coli</i>	Meat	<i>blaSHV-5, blaCMY-13, blaVIM-1</i>	No/No	Belgium	(Garcia-Graells et al. 2020)
<i>E. coli, K. pneumoniae</i>	Chicken, pork, beef,	<i>blaSHV; blaTEM; blaCTX-M; blaCMY; blaOXA</i>	Yes ⁴ /Yes	Burma	(Sugawara et al. 2019)
<i>E. coli, K. pneumoniae</i>	Pork, chicken, and beef	<i>blaNDM-1,-5,-7</i>	No/Yes	China	(Zhang et al. 2019)
<i>E. coli, K. pneumoniae</i>	Raw beef	<i>blaSHV; blaTEM; blaCTX-M; blaCMY-1,-2; blaOXA</i>	No/No	South Africa	(Montso et al. 2019)
<i>E. coli, K. pneumoniae</i>	Chicken meat	<i>blaSHV-2,-5,-12,-27; blaTEM-15,-52B,-52C; blaCTX-M-1,-2,-15,-32</i>	No/Yes	Netherlands	(Huizinga et al. 2019)
<i>E. coli</i>	Raw chicken meat	<i>blaSHV-12; blaTEM-1b; blaCTX-M-1,-55</i>	Yes/No	Tunisia	(Hassen et al. 2020)
<i>E. coli</i>	Chicken meat	<i>blaCTX-M-2,-8,-15; blaCMY-2</i>	Yes/No	Brazil	(Botelho et al. 2020)
<i>E. coli</i>	Chicken, turkey, beef and pork	<i>blaTEM-1b,-1D; blaCTX-M-1,-14,-15,-27,-32; blaHERA-3</i>	No/Yes	United States	(Tadesse et al. 2018)
<i>E. coli</i>	Chicken, beef, pork	<i>blaSHV, blaTEM, blaCTX-M</i>	No/No	China	(Ye et al. 2018)
<i>E. coli</i>	Chicken and pork	<i>blaSHV-12; blaTEM-176; blaCTX-M-1,-2,-3,-8,-14,-25,-55,-65</i>	No/No	Japan	(Hayashi et al. 2018)
<i>E. coli</i>	Beef	<i>blaTEM; blaCTX-M-1</i>	No/No	Portugal	(Clemente et al. 2019)
<i>E. coli</i>	Beef	<i>blaCTX-M-1,-3,-14,-15,-24,-32</i>	Yes/No	Alegria	(Rebbah et al. 2018)

Resistant Bacteria in Food

<i>E. coli</i>	Pork	<i>blaTEM-1; blaCTX-M-1,-14,-15; blaCMY-2</i>	<u>No</u> /No	Germany	(Schill et al. 2017)
<i>E. coli</i>	Pork	<i>blaCTX-M-14,-27,-55,-65</i>	<u>Yes</u> ⁴ /No	Vietnam	(Hoang et al. 2017)
<i>E. coli</i>	Beef, chicken, pork	<i>blaSHV-2,-2a,-12; blaTEM-1b,-52,-52c; blaCTX-M-1,-2,-32,-55</i>	<u>No</u> /No	United Kingdom	(Randall et al. 2017)
<i>E. coli</i>	Pork	<i>blaSHV; blaTEM; blaCTX-M-1,-9</i>	<u>No</u> /No	China	(Li et al. 2016)
<i>E. coli</i>	Chicken meat and cow milk	<i>blaSHV; blaTEM; blaCTX-M-1,-8</i>	<u>No</u> /No	Turkey	(Tekiner and Özpınar 2016)
<i>E. coli</i>	Chicken meat	<i>blaSHV-12; blaTEM-1B; blaCTX-M-1</i>	<u>No</u> /Yes	Netherlands	(Kluytmans-van den bergh et al. 2015)
<i>E. coli</i>	Chicken meat	<i>blaSHV; blaTEM; blaCTX-M-1,-15,-61</i>	<u>No</u> /No	Ghana	(Rasmussen et al. 2015)
<i>E. coli, K. pneumoniae</i>	Chicken meat	<i>blaSHV; blaTEM; blaCTX-M</i>	<u>No</u> /No	Egypt	(Abdallah et al. 2015)
<i>E. coli</i>	Chicken and beef	<i>blaCTX-M-1, blaSHV-5,-12; blaTEM-1b; blaCTX-M-1,-3,-15; blaCMY-2</i>	<u>No</u> /No	Turkey	(Pehlivanlar Önen et al. 2015)
<i>E. coli</i>	Pork and beef	<i>blaCTX-M, blaTEM, blaSHV, blaCMY</i>	<u>No</u> /No	Germany	(Kaesbohrer et al. 2019)

Vegetable based foods

<i>E. coli</i>	Vegetables ¹	<i>blaNDM-5,-9; mcr-1</i>	<u>Yes</u> /No	China	(B. T. Liu and Song 2019)
<i>E. coli, K. pneumoniae</i>	Vegetables ¹	<i>blaCTX-M-1G; blaNDM-5; blaKPC-2</i>	<u>Yes</u> /No	China	(B. T. Liu et al. 2018)
<i>E. coli</i>	Vegetables	<i>blaCTX-M</i>	<u>No</u> /No	Germany	(Kaesbohrer et al. 2019)
<i>K. pneumoniae</i>	Lettuce	<i>blaSHV, blaTEM</i>	<u>Yes</u> /No	United States	(Bhutani et al. 2015)

Resistant Bacteria in Food

<i>E. coli, K. pneumoniae</i>	Vegetables ¹	<i>blaSHV-1,-2,-11,-12,-27,-28,-61; blaTEM-1; blaCTX-M-14,-15,-55</i>	<u>No</u> /No	Korea	(Kim et al. 2015)
<i>K. pneumoniae</i>	Vegetables ¹	<i>blaSHV-12, blaCTX-M-14</i>	<u>No</u> /No	Dutch	(Reuland et al. 2014)
<i>E. coli, K. pneumoniae</i>	Vegetables ¹	<i>blaSHV; blaTEM; blaCTX-M; blaCMY; blaDHA; blaOXA</i>	<u>Yes</u> ⁴ /Yes	Burma	(Sugawara et al. 2019)

Dairy based food

<i>E. coli, K. pneumoniae</i>	Cow milk	<i>blaCTX-M-15</i>	<u>No</u> /No	Tunisia	(Saidani et al. 2018)
-------------------------------	----------	--------------------	---------------	---------	-----------------------

¹ Includes a variety of vegetables reported in the selected publication, for details see the references.


² Only resistance genes conferring resistance to carbapenems, cephalosporins, and colistin are listed. Original articles might report resistance genes to other antibiotics.

³ In the case that more than one variants on a given resistance gene were reported, these are listed as this example: *blaNDM-1,-5,-7* for *blaNDM-1*, *blaNDM-5*, *blaNDM-7*


⁴ PFGE was used for plasmid analysis instead of for molecular typing

Figure 1.JPEG

A. Sample origins



B. Bacterial isolation and data collection



C. Expected results

